

GAMMAGARD *LIQUID*

[Immune Globulin Infusion (Human)] 10%

TAKE A STEP WITH GAMMAGARD LIQUID

A therapy to improve neuromuscular disability and impairment in **adult patients with CIDP**.

Talk to your doctor to find out if **GAMMAGARD LIQUID** is right for you.



What is GAMMAGARD LIQUID?

GAMMAGARD LIQUID is a ready-to-use liquid medicine that is given in a vein (intravenously) to treat chronic inflammatory demyelinating polyneuropathy (CIDP) in adults.

IMPORTANT SAFETY INFORMATION

What is the most important information that I should know about GAMMAGARD LIQUID?

GAMMAGARD LIQUID can cause the following serious reactions:

- Severe allergic reactions causing difficulty in breathing or skin rashes
- Decreased kidney function or kidney failure
- Blood clots in the heart, brain, lungs or elsewhere in the body
- Severe headache, drowsiness, fever, painful eye movements, or nausea and vomiting
- Dark colored urine, swelling, fatigue, or difficulty breathing

Please see additional Important Safety Information throughout, [click here for accompanying Information For Patients](#), and discuss with your HCP.



About CIDP

With CIDP, you may experience a range of symptoms over time.

CIDP is a rare autoimmune disease which targets the nerves, leading to muscle weakness and numbness. IgG is thought to reduce the damage to the nerve and assist in defending the nerve from harm.

Over time, the following symptoms may come and go, but it's also not uncommon for them to get worse:



**Arm/leg
weakness**



**Difficulty with
balance**



**Difficulty with fine
motor skills**



**Loss of
coordination**



**Walking aid
usage**

**CIDP can cause arm and leg impairment.
Examples of activities that can be affected may include:**



**Handling
small coins**



**Brushing
hair**



**Turning a key
in a lock**



**Working
a zipper
or buttons**



**Washing
hair**

Treatment for CIDP

Finding a treatment plan for managing your CIDP will require working closely with your healthcare team. EFNS/PNS guidelines strongly recommend IVIG as a treatment option for CIDP.



Gammagard Liquid Clinical Trial Results

In a clinical study with CIDP patients who have relapsed, 94% of people in the clinical study (17 out of 18 patients) experienced improvement in a tool that measures arm and leg disability after 6 months of treatment.



Reduced disability



**Improvement of
limb impairment**

IMPORTANT SAFETY INFORMATION (continued)

Who should not use GAMMAGARD LIQUID?

Do not use GAMMAGARD LIQUID if you:

- Have had a severe allergic reaction to immune globulin or other blood products.
- Have a condition called selective (or severe) immunoglobulin A (IgA) deficiency.



The most common side effects of GAMMAGARD LIQUID for CIDP include:

- Headache
- Fever
- Anemia
- Illness
- Dizziness
- Migraine
- Drowsiness
- Tremor
- Nasal dryness
- Upper abdominal pain
- Vomiting
- Chills
- Head cold
- Pain in extremity

These are not all the possible side effects. Talk to your healthcare provider about any side effect that bothers you or that does not go away.

IMPORTANT SAFETY INFORMATION (continued)

What should I avoid while taking GAMMAGARD LIQUID?

- GAMMAGARD LIQUID can make vaccines (like measles/mumps/rubella or chickenpox vaccines) not work as well for you. Before you get any vaccines, tell your healthcare provider (HCP) that you take GAMMAGARD LIQUID.
- Tell your HCP if you are pregnant, or plan to become pregnant, or if you are nursing.

Please see additional Important Safety Information throughout and click for accompanying [Information For Patients](#), including Warning about Blood Clots, Decreased Kidney Function, and Kidney Failure, and discuss with your HCP.

GAMMAGARD LIQUID

[Immune Globulin Infusion (Human)] 10%

Beginning Your Treatment Plan

GAMMAGARD LIQUID is an immune globulin infusion therapy to improve neuromuscular disability and impairment in adult patients with CIDP. It is a ready-to-use liquid that is given in a vein (intravenously) to help adult patients manage some CIDP symptoms.



Care options

You can receive your GAMMAGARD LIQUID treatments at an **infusion center**, in your **doctor's office** or another **medical setting**, or at **home with an infusion nurse**.



Induction Dosing

Your doctor will decide when you will receive your induction doses over 2 to 5 consecutive days.



Maintenance Dosing

Your doctor will then plan for your maintenance dosing, usually over 1 to 4 consecutive days, every 3 weeks. This may be adjusted based on your treatment response.

IMPORTANT SAFETY INFORMATION (continued)

What are the possible or reasonably likely side effects of GAMMAGARD LIQUID?

GAMMAGARD LIQUID can cause serious side effects. If any of the following problems occur after starting GAMMAGARD LIQUID, stop the infusion immediately and contact your HCP or call emergency services:

- Hives, swelling in the mouth or throat, itching, trouble breathing, wheezing, fainting or dizziness. These could be signs of a serious allergic reaction.
- Bad headache with nausea, vomiting, stiff neck, fever, and sensitivity to light. These could be signs of irritation and swelling of the lining around your brain.
- Reduced urination, sudden weight gain, or swelling in your legs. These could be signs of a kidney problem.
- Pain, swelling, warmth, redness, or a lump in your legs or arms. These could be signs of a blood clot.

Infusion Center Information

An infusion center is an outpatient clinic that is certified to administer infusion therapy. It is often a cozy setting staffed with knowledgeable people who specialize in infusion therapy. These facilities have advanced medical equipment and often provide personalized care to meet your needs.

Some people use their infusion time to nap, read a book, or phone a friend. Consider what you might want to do during your infusion time. Tell your nurse if you need anything that will make you more comfortable. Often, nurses can dim the lights, bring you a blanket, or offer a heat pack for your IV site.

There are many individuals involved in a single infusion treatment, all working together to ensure a safe and comfortable experience. From the prescribing doctor, to the facility staff, the administrators, and the infusion nurses, each role contributes to patient care in an important way.

Additional Support

Here are some questions that can help you start important conversations about your treatment with **your healthcare provider**.

- **How can I expect my symptoms to change over time?**
- **What impact will CIDP have on my daily life?**
- **How does my CIDP management plan fit into my lifestyle?**
- **What will IVIG feel like during the infusion?**
- **What side effects are common?**
- **When should I contact an HCP after my infusion?**
- **What are my next steps for treatment?**

IMPORTANT SAFETY INFORMATION (continued)

- Brown or red urine, fast heart rate, yellow skin or eyes. These could be signs of a liver or blood problem.
- Chest pain or trouble breathing, or blue lips or extremities. These could be signs of a serious heart or lung problem.
- Fever over 100°F. This could be a sign of an infection.

Please see additional Important Safety Information throughout and click for accompanying [Information For Patients](#), including Warning about Blood Clots, Decreased Kidney Function, and Kidney Failure, and discuss with your HCP.

Takeda Patient Support

When you're prescribed GAMMAGARD LIQUID, Takeda Patient Support is here for you. Our support specialists can address your questions and concerns and help get you the answers, resources, and tools you need. Some of the ways we can assist include:



Working with your specialty pharmacy (or site of care) to help you receive your prescribed Takeda treatment



Connecting you to co-pay or insurance help



Arranging for nursing support if you have questions about your condition and your prescribed Takeda treatment

VISIT:
TAKEDAPATIENTSUPPORT.COM

Depending on where you are in your treatment journey, Takeda also has another treatment option for CIDP, so it is important to **talk to your doctor about the treatment options that may be right for you.**

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of GAMMAGARD LIQUID for CIDP include:

- | | | | |
|------------|--------------|------------------------|---------------------|
| • Headache | • Dizziness | • Nasal dryness | • Chills |
| • Fever | • Migraine | • Upper abdominal pain | • Nasopharyngitis |
| • Anemia | • Somnolence | | • Pain in extremity |
| • Illness | • Tremor | • Vomiting | |

Please see additional Important Safety Information throughout and click for accompanying [Information For Patients](#), including Warning about Blood Clots, Decreased Kidney Function, and Kidney Failure, and discuss with your HCP.

Resources

CIDP Advocacy Groups

There are many helpful websites for those living with CIDP and their caregivers, which you can turn to for more information and support.

The GBS/CIDP Foundation International

The GBS/CIDP Foundation International is a nonprofit organization that supports and advocates for people with CIDP.



American Association of Neuromuscular & Electrodiagnostic Medicine

The American Association of Neuromuscular & Electrodiagnostic Medicine is a nonprofit association dedicated to improving the quality of medical care provided to people with muscle and nerve disorders.

American Neuromuscular Foundation (ANF)

A nonprofit organization dedicated to strengthening the global effort to cure neuromuscular disease. See the work and programs for people with neuromuscular diseases.

Caregiver Action Network (CAN)

Caregiver Action Network is an organization that works to improve the quality of life for the millions of Americans who care for loved ones with chronic conditions, disabilities, disease, or the frailties of old age.

The Foundation for Peripheral Neuropathy

The Foundation for Peripheral Neuropathy is a public charity foundation that aims to improve the lives of people living with neuropathy through education, awareness, and research. This website includes information on finding a support group (in-person and online group options included).

Neuropathy Action Foundation (NAF)

An advocacy organization with a virtual Resource Center with information on CIDP, as well as stories from people living with the disorder.

What is GAMMAGARD LIQUID?

GAMMAGARD LIQUID is a ready-to-use liquid medicine that is given in a vein (intravenously) to treat chronic inflammatory demyelinating polyneuropathy (CIDP) in adults.

IMPORTANT SAFETY INFORMATION

What is the most important information that I should know about GAMMAGARD LIQUID?

GAMMAGARD LIQUID can cause the following serious reactions:

- Severe allergic reactions causing difficulty in breathing or skin rashes
- Decreased kidney function or kidney failure
- Blood clots in the heart, brain, lungs or elsewhere in the body
- Severe headache, drowsiness, fever, painful eye movements, or nausea and vomiting
- Dark colored urine, swelling, fatigue, or difficulty breathing

Who should not use GAMMAGARD LIQUID?

Do not use GAMMAGARD LIQUID if you:

- Have had a severe allergic reaction to immune globulin or other blood products.
- Have a condition called selective (or severe) immunoglobulin A (IgA) deficiency.

What should I avoid while taking GAMMAGARD LIQUID?

- GAMMAGARD LIQUID can make vaccines (like measles/mumps/rubella or chickenpox vaccines) not work as well for you. Before you get any vaccines, tell your healthcare provider (HCP) that you take GAMMAGARD LIQUID.
- Tell your HCP if you are pregnant, or plan to become pregnant, or if you are nursing.

What are the possible or reasonably likely side effects of GAMMAGARD LIQUID?

GAMMAGARD LIQUID can cause serious side effects. If any of the following problems occur after starting GAMMAGARD LIQUID, stop the infusion immediately and contact your HCP or call emergency services:

- Hives, swelling in the mouth or throat, itching, trouble breathing, wheezing, fainting or dizziness. These could be signs of a serious allergic reaction.
- Bad headache with nausea, vomiting, stiff neck, fever, and sensitivity to light. These could be signs of irritation and swelling of the lining around your brain.
- Reduced urination, sudden weight gain, or swelling in your legs. These could be signs of a kidney problem.
- Pain, swelling, warmth, redness, or a lump in your legs or arms. These could be signs of a blood clot.

IMPORTANT SAFETY INFORMATION (CONTINUED)

What are the possible or reasonably likely side effects of GAMMAGARD LIQUID?

GAMMAGARD LIQUID can cause serious side effects. If any of the following problems occur after starting GAMMAGARD LIQUID, stop the infusion immediately and contact your HCP or call emergency services:

- Brown or red urine, fast heart rate, yellow skin or eyes. These could be signs of a liver or blood problem.
- Chest pain or trouble breathing, or blue lips or extremities. These could be signs of a serious heart or lung problem.
- Fever over 100°F. This could be a sign of an infection.

The most common side effects of GAMMAGARD LIQUID for CIDP include:

- | | |
|--------------|------------------------|
| • Headache | • Tremor |
| • Fever | • Nasal dryness |
| • Anemia | • Upper abdominal pain |
| • Illness | • Vomiting |
| • Dizziness | • Chills |
| • Migraine | • Nasopharyngitis |
| • Somnolence | • Pain in extremity |

These are not all the possible side effects. Talk to your HCP about any side effect that bothers you or that does not go away.

For additional safety information, click for [Information For Patients](#) and discuss with your HCP.

You are encouraged to report negative side effects of prescription drugs to FDA. Visit www.fda.gov/medwatch, or call [1-800-FDA-1088](tel:1-800-FDA-1088).



For information on managing CIDP with GAMMAGARD,

VISIT: WWW.GAMMAGARD.COM/CIDP

IMPORTANT SAFETY INFORMATION

What is the most important information that I should know about GAMMAGARD LIQUID?

GAMMAGARD LIQUID can cause the following serious reactions:

- Severe allergic reactions causing difficulty in breathing or skin rashes
- Decreased kidney function or kidney failure
- Blood clots in the heart, brain, lungs or elsewhere in the body
- Severe headache, drowsiness, fever, painful eye movements, or nausea and vomiting
- Dark colored urine, swelling, fatigue, or difficulty breathing

Please see additional Important Safety Information throughout and click here for accompanying [Information For Patients](#), including Warning about Blood Clots, Decreased Kidney Function, and Kidney Failure, and discuss with your HCP.

©2024 Takeda Pharmaceuticals U.S.A., Inc., 500 Kendall Street, Cambridge, MA 02142.
1-877-TAKEDA-7 (1-877-825-3327). All rights reserved. TAKEDA®, the TAKEDA Logo®, and the TAKEDA Patient Support Logo® are registered trademarks of Takeda Pharmaceutical Company Limited. GAMMAGARD LIQUID® is a registered trademark of Baxalta Incorporated.
US-GGL-0624v1.0 08/24

