



GAMMAGARD LIQUID Patient Start Form

Fax pages 1-4 to 1-866-861-1752 | Phone: 1-866-861-1750

Please ensure patient reads and signs pages 3 and 4 for appropriate authorizations.



[Immune Globulin Infusion (Human)] 10%

1 Prescribing Physician Information

Prescriber Name (First, Last): _____ Title: _____

State License #: _____ NPI #: _____ Tax ID #: _____ PTAN #: _____

Street Address: _____ City: _____ State: _____ Zip Code: _____

Office Contact: _____ Email: _____

Telephone: _____ Fax: _____

2 Patient Information

Male Female

Patient Name (First, Middle Initial, Last): _____

DOB (MM/DD/YYYY): _____ Last 4 Digits of Social Security #: _____ Email: _____

Street Address: _____

City: _____ State: _____ Zip Code: _____

Mobile Telephone: _____ Home Telephone: _____

Caregiver Name (First, Last): _____ Relationship to Patient: _____

Caregiver Telephone: _____ Caregiver Email: _____

3 Insurance Information

Please attach copies of both sides of patient's medical and prescription insurance cards.

Check if patient does not have insurance.

Primary Insurance: _____	Pharmacy Plan Name: _____	Secondary Insurance: _____
Insurance Telephone: _____	Pharmacy Plan Telephone: _____	Insurance Telephone: _____
Policy ID #: _____	Policy ID #: _____	Policy ID #: _____
Group ID #: _____	Group ID #: _____	Group ID #: _____
Policy Holder Name: _____	Rx BIN #: _____	Policy Holder Name: _____
Policy Holder DOB (MM/DD/YYYY): _____	Rx PCN #: _____	Policy Holder DOB (MM/DD/YYYY): _____

Patient Name: _____

Fax pages 1-4 to 1-866-861-1752 | Phone: 1-866-861-1750

4 Diagnosis/Medical Assessment

Diagnosis (ICD-10): _____

PI _____

IgA Level (mg/dL): _____

IgG Level (mg/dL): _____

IgM Level (mg/dL): _____

Pre-Titer Level (mcg/mL): _____

Post-Titer Level (mcg/mL): _____

CIDP _____

EMG/NCS: _____

Optional MRI: _____

Optional CSF: _____

Other: _____

MMN _____

MRI Results: _____

Optional Anti-GM1 Titer (mg/mL): _____

EMG/NCS/Nerve Ultrasound (m/sec): _____

Other: _____

5 GAMMAGARD LIQUID Prescription, Training Request/Waiver, and Prescribing Physician Signature

Name (First, Middle Initial, Last): _____

DOB (MM/DD/YYYY): _____

Patient Weight (kg): _____

Prescription: GAMMAGARD LIQUID® [Immune Globulin Infusion (Human)] 10% Solution

- Intravenous immune globulin (IVIG) administration** (for PI, MMN, and CIDP)¹
- For patients with primary immunodeficiency (PI), administer IVIG doses of 300 to 600 mg/kg every 3 or 4 weeks based on clinical response. See the Infusion Rates for IV Administration table on page 5 for calculation of infusion rate.
- For patients with multifocal motor neuropathy (MMN), administer IVIG doses of 0.5 to 2.4 grams/kg per month based on clinical response.
- For patients with chronic inflammatory demyelinating polyneuropathy (CIDP), induction dose is 2 g/kg in divided doses over 2 to 5 consecutive days, followed by maintenance infusions. The maintenance dose is 1 g/kg in divided doses over 1 to 4 consecutive days, every 3 weeks. The maintenance dose and dosing interval may be adjusted according to clinical response.

- Subcutaneous immune globulin (SCIG) administration** (for PI only)¹
- For patients with PI switching from IVIG to SCIG treatment, the formula below is used to calculate the recommended initial dose. See the Infusion Rates for SC Administration table on page 5 for calculation of infusion rate. Maintenance dose is based on clinical response and target IgG trough level.

To calculate SCIG dose = (1.37 x previous IVIG dose) ÷ Number of weeks between IVIG doses

Special Instructions: _____

Patient is already on GAMMAGARD LIQUID

Refills (as allowed by state or payer requirement)

Ordered Dose (grams): _____

Every (weeks): _____

Route: _____

Central IV Peripheral IV SC needle

Length (mm): _____

No known drug allergies

Patient allergies (drug and non-drug): _____

Preferred site of care if not self-administered (check one) _____

- Infusion suite Begin treatment in clinical setting, then transition to home care

Preferred Specialty Pharmacy: _____

Has a referral been sent to site of care? Yes No

- Prescriber's office Home infusion Hospital outpatient

Preferred Infusion Suite/Hospital Outpatient (if applicable): _____

Additional services and infusion training are available.

Check the box next to any of the services below. Check the last box if the patient opts out.

- Pharmacy to provide needles, syringes, venous access device supplies, and other ancillary supplies needed for infusion
- Durable medical equipment (DME)—infusion pump with supplies
- Name of pharmacy (to provide anaphylactic kit) _____

Training available to SCIG patients

GAMMAGARD LIQUID SCIG is intended for self-administration or administration by a caregiver. The patient or caregiver should be trained by a healthcare professional. Takeda Patient Support provides free infusion training services to enrolled patients.

If you choose to opt out of these services, please check this box.

By signing this form, I certify that therapy with GAMMAGARD LIQUID is medically necessary for the patient identified in this application ("Patient"). I have reviewed the current GAMMAGARD LIQUID Prescribing Information and will be supervising Patient's treatment. I have received from Patient, or his/her personal representative, the necessary authorization to release, in accordance with applicable federal and state law regulations, referenced medical and/or other patient information relating to GAMMAGARD LIQUID therapy to Takeda Pharmaceutical Company Limited, including its agents or contractors, for the purpose of seeking information related to coverage and/or assisting in initiating or continuing GAMMAGARD LIQUID therapy. I authorize Takeda Patient Support to transmit this prescription to the appropriate pharmacy designated by me, Patient, or Patient's plan. I agree that product provided through the Program shall only be used for Patient, must not be resold, offered for sale or trade or returned for credit.

Prescriber Signature (Required) Stamps not acceptable

SIGN

DISPENSE AS WRITTEN _____ Date _____ SUBSTITUTION PERMITTED _____ Date _____

The prescriber is required to comply with his/her state-specific prescription requirements such as e-prescribing, state-specific prescription form, fax language, etc. Non-compliance with state-specific requirements could result in delay.

Please see Important Safety Information on page 6 and click for Full Prescribing Information, including Boxed Warning regarding Thrombosis, Renal Dysfunction and Acute Renal Failure.

Patient Name: _____

Fax pages 1-4 to 1-866-861-1752 | Phone: 1-866-861-1750

6 Patient HIPAA Authorization

Patient Name (First, Middle Initial, Last): _____

DOB (MM/DD/YYYY): _____

By signing the Patient Authorization section on the third page of this Takeda Patient Support Ig Enrollment Form, I authorize my physician, health insurance, and pharmacy providers (including any specialty pharmacy that receives my prescription) to disclose my protected health information, including, but not limited to, information relating to my medical condition, treatment, care management, and health insurance, as well as all information provided on this form (“Protected Health Information”), to Takeda Pharmaceuticals U.S.A., Inc. and its present or future affiliates, including the affiliates and service providers that work on Takeda’s behalf in connection with the Takeda Patient Support, Ig Patient Support Program (the “Companies”). The Companies will use my Protected Health Information for the purpose of facilitating the provision of the Takeda Patient Support, Ig Patient Support Program products, supplies, or services as selected by me or my physician and may include (but not be limited to) verification of insurance benefits and drug coverage, prior authorization education, financial assistance with co-pays, patient assistance programs, and other related programs. Specifically, I authorize the Companies to 1) receive, use, and disclose my Protected Health Information in order to enroll me in Takeda Patient Support, Ig and contact me, and/or the person legally authorized to sign on my behalf, about Takeda Patient Support, Ig; 2) provide me, and/or the person legally authorized to sign on my behalf, with educational materials, information, and services related to Takeda Patient Support, Ig; 3) verify, investigate, and provide information about my coverage for GAMMAGARD LIQUID, including but not limited to communicating with my insurer, specialty pharmacies, and others involved in processing my pharmacy claims to verify my coverage; 4) coordinate prescription fulfillment; and 5) use my information to conduct internal analyses. I understand that employees of the Companies only use my Protected Health Information for the purposes described herein, to administer the Takeda Patient Support, Ig Patient Support Program or as otherwise required or allowed under the law, unless information that specifically identifies me is removed. Further, I understand that my physician, health insurance, and pharmacy providers may receive financial remuneration from the Companies for providing Protected Health Information, which may be used for marketing purposes. I understand that Protected Health Information disclosed under this Authorization may no longer be protected by federal privacy law. I understand that I am entitled to a copy of this Authorization. I understand that I may revoke this Authorization and that instructions for doing so are contained in Takeda’s Website Privacy Notice available at www.takeda.com/privacy-notice/ or I may revoke this Authorization at any time by sending written notice of revocation to Takeda Patient Services 610 Crescent Executive Court, Suite 200 Lake Mary, FL 32746. I understand that such revocation will not apply to any information already used or disclosed through this Authorization. This Authorization will expire within five (5) years from the date it is signed and provided on the first page of this enrollment form, unless a shorter period is provided for by state law. I understand that I may refuse to sign this Authorization and that refusing to sign this Authorization will not change the way my physician, health insurance, and pharmacy providers treat me. I also understand that if I do not sign this Authorization, I will not be able to receive Takeda Patient Support, Ig Patient Support Program products, supplies, or services.

SIGN

Signature of Patient (Required) _____

Date _____

SIGN

*Legal Representative Signature _____

Date _____

*Legal Representative Name: _____

*Relationship to Patient: _____

*Required only if applicable.

Please see Important Safety Information on page 6 and click for Full Prescribing Information, including Boxed Warning regarding Thrombosis, Renal Dysfunction and Acute Renal Failure.

Patient Name: _____

Fax pages 1-4 to 1-866-861-1752 | Phone: 1-866-861-1750

7 Takeda Patient Support Enrollment (signature required for enrollment)

By signing below, I am electing to enroll in Takeda Patient Support Services (“Services”) and direct all disclosures of my Information in connection with such Services (which may include, but are not limited to, verification of insurance benefits and drug coverage, prior authorization support, financial assistance with co-pays, patient assistance programs, alternate funding sources, other related programs, communication with me or my prescribing physician by mail, email, or telephone about my medical condition, treatment, care management, product information, and health insurance).

SIGN 

Signature of Patient (Required)/*Legal Representative Signature

Date

*Required only if applicable.

8 Patient Consent for Communications

Marketing Communications

- By checking this box, I authorize the use of my Information for Takeda marketing activities and consent to receiving marketing, market research opportunities, and promotional communications from Takeda. I hereby give consent to Takeda, its affiliates, and their agents and representatives to send communications and information to me via the contact information I have provided above. I understand that this consent will be in effect until I cancel such authorization.

Text Communications

- By checking this box, you agree to the Takeda Patient Support (“Program”) text message terms and conditions below, and you agree to receive text messages on your mobile device subject to the Terms & Conditions. You consent to receive autodialed and/or prerecorded calls and/or text messages from or on behalf of the Program at the telephone number provided above. You understand that this consent is not a condition of purchase or use of the Program or of any Takeda product or service. Such messages may be nonmarketing messages related to the Patient Support Program. There is no fee payable to Takeda to receive text messages; however, your carrier’s message and data rates may apply.

Text Communication Agreement Terms & Conditions

You represent that you are the account holder for the mobile telephone number(s) that you provide to opt in to the Program. You are responsible for notifying Takeda immediately if you change your mobile telephone number. You may notify Takeda of a number change by calling 1-866-861-1750. Data obtained from you in connection with your registration for, and use of, this SMS service may include your phone number and/or email address, related carrier information, and elements of pharmacy claim information and will be used to administer this Program and to provide Program benefits such as information about your prescription, refill reminders, and Program updates and alerts.

Takeda will not be liable for any delays in the receipt of any SMS messages, as delivery is subject to effective transmission from your network operator. This Program is valid with most major US cellular providers.

Takeda may be required to contact the user if an adverse event is reported.

You agree to indemnify Takeda and any third parties texting on its behalf in full for all claims, expenses, and damages related to or caused, in whole or in part, by your failure to immediately notify us if you change your telephone number, including but not limited to all claims, expenses, and damages related to or arising under the Telephone Consumer Protection Act.

Takeda reserves the right to rescind, revoke, or amend the Program without notice at any time.

You can unsubscribe from this Program by texting back STOP to any message or by calling 1-866-861-1750.

Patient Name:

Fax pages 1-4 to **1-866-861-1752** | Phone: **1-866-861-1750**

Before you fax this form, confirm you have:

- Completed **sections 1-8** Attached a copy of the patient's **insurance card (front and back)** Ensured the patient/legal representative has **signed pages 3 and 4** Not submitted any documentation of labs, clinical history, or other documents supporting the prior authorization process

- 1 Prescribing Physician Information** **2 Patient Information** **3 Insurance Information** **4 Diagnosis/Medical Assessment**

5 GAMMAGARD LIQUID Prescription, Training Request/Waiver, and Prescribing Physician Signature

- Please indicate the number of refills • This is a prescription; a physician's signature and date are required
- **Available to SCIG patients only:** Check the appropriate box to specify whether you would like your patient to be trained by Takeda on self-administration or whether training has already occurred

	Infusion Rates for IV Administration (PI) ¹	Infusion Rates for IV Administration (MMN) ¹	Infusion Rates for IV Administration (CIDP) ¹
Initial	0.5 mL/kg/hr (0.8 mg/kg/min) for 30 minutes	0.5 mL/kg/hr (0.8 mg/kg/min)	0.5 mL/kg/hr (0.8 mg/kg/min)
Subsequent	Increase every 30 minutes (if tolerated) up to 5 mL/kg/hr (8 mg/kg/min)	Infusion rate may be increased if tolerated up to 5.4 mL/kg/hr (9 mg/kg/min)	Infusion rate may be increased if tolerated up to 5.4 mL/kg/hr (9 mg/kg/min)

	Infusion Rates for SC Administration (PI) ¹	
	Patients ≥40 kg	Patients <40 kg
Initial	30 mL/site at a rate of 20 mL/hr/site	20 mL/site at a rate of 15 mL/hr/site
Maintenance	30 mL/site at a rate of 20 to 30 mL/hr/site	20 mL/site at a rate of 15 to 20 mL/hr/site

6 Patient HIPAA Authorization

The patient signature is required to allow personal health information to be shared by third parties to Takeda to facilitate access to GAMMAGARD LIQUID (insurance benefits, self-administration training [available to SCIG patients only], transfer Rx to specialty pharmacy provider, etc).

7 Takeda Patient Support Enrollment

The patient signature is required to enroll in Takeda Patient Support, and allows patients to receive product support and services from Takeda if eligible. Check additional services and infusion training if needed, or check the last box if patient opts out.

8 Patient Consent for Communications

The patient must check the appropriate boxes to receive Takeda marketing and text communications.

What happens next?

- Once the completed form has been submitted to Takeda Patient Support, a dedicated Support Specialist will be assigned to your eligible patient
- The Support Specialist will contact the patient directly to inform him or her of the services available through Takeda Patient Support and to begin the insurance verification process
- The Support Specialist will work with the insurance company to determine insurance benefits
- The Support Specialist will assess the patient's eligibility for co-pay support (when applicable) and provide information about third-party financial assistance programs, as necessary
- **Available to SCIG patients only:** If requested, the Support Specialist will set up Takeda-provided self-administration training services

INDICATIONS

GAMMAGARD LIQUID is indicated as a replacement therapy for primary humoral immunodeficiency (PI) in adult and pediatric patients two years of age or older, as a maintenance therapy to improve muscle strength and disability in adult patients with Multifocal Motor Neuropathy (MMN), and as a therapy to improve neuromuscular disability and impairment in adult patients with Chronic Inflammatory Demyelinating Polyneuropathy (CIDP).

LIMITATIONS OF USE (CIDP): GAMMAGARD LIQUID has not been studied in immunoglobulin-naïve patients with CIDP. GAMMAGARD LIQUID maintenance therapy in CIDP has not been studied for periods longer than 6 months. After responding during an initial treatment period, not all patients require indefinite maintenance therapy with GAMMAGARD LIQUID in order to remain free of CIDP symptoms. Individualize the duration of any treatment beyond 6 months based upon the patient's response and demonstrated need for continued therapy.

GAMMAGARD LIQUID for PI is for intravenous or subcutaneous use.
GAMMAGARD LIQUID for MMN and CIDP is for intravenous use only.

IMPORTANT SAFETY INFORMATION

WARNING: THROMBOSIS, RENAL DYSFUNCTION, and ACUTE RENAL FAILURE

- **Thrombosis may occur with immune globulin (IG) products, including GAMMAGARD LIQUID. Risk factors may include advanced age, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of estrogens, indwelling vascular catheters, hyperviscosity, and cardiovascular risk factors. Thrombosis may occur in the absence of known risk factors.**
- **Renal dysfunction, acute renal failure, osmotic nephrosis, and death may occur in predisposed patients with immune globulin intravenous (IGIV) products. Patients predisposed to renal dysfunction include those with any degree of pre-existing renal insufficiency, diabetes mellitus, age greater than 65, volume depletion, sepsis, paraproteinemia, or patients receiving known nephrotoxic drugs. Renal dysfunction and acute renal failure occur more commonly in patients receiving IGIV products containing sucrose. GAMMAGARD LIQUID does not contain sucrose.**
- **For patients at risk of thrombosis, administer GAMMAGARD LIQUID at the minimum dose and infusion rate practicable. Ensure adequate hydration in patients before administration. Monitor for signs and symptoms of thrombosis and assess blood viscosity in patients at risk of hyperviscosity.**

Contraindications

- History of anaphylactic or severe systemic hypersensitivity reactions to human IG.
- IgA-deficient patients with antibodies to IgA and a history of hypersensitivity to human IG. Anaphylaxis has been reported with intravenous (IV) use of GAMMAGARD LIQUID.

Warnings and Precautions

Hypersensitivity: Severe hypersensitivity reactions may occur, even in patients who have tolerated previous treatment with human IG. If a hypersensitivity reaction occurs, discontinue infusion immediately and institute appropriate treatment. IgA-deficient patients with antibodies to IgA are at greater risk of developing potentially severe hypersensitivity reactions, including anaphylaxis.

Renal Dysfunction/Failure: Acute renal dysfunction/failure, acute tubular necrosis, proximal tubular nephropathy, osmotic nephrosis, and death may occur with IV use of IG products, especially those containing sucrose. Acute renal dysfunction/failure has been reported in association with infusions of GAMMAGARD LIQUID. Ensure patients are not volume depleted prior to infusion. In patients at risk due to pre-existing renal insufficiency or predisposition to acute renal failure, assess renal function before initiation and throughout treatment, and use the minimum infusion rate practicable for IV administration. If renal function deteriorates, consider discontinuation.

Hyperproteinemia, increased serum viscosity, and hyponatremia may occur. It is critical to distinguish true hyponatremia from a pseudohyponatremia because certain treatments may lead to volume depletion, a further increase in serum viscosity, and a predisposition to thromboembolic events.

Thrombosis: May occur following treatment with IG products and in the absence of known risk factors. In patients at risk, administer at the minimum dose and infusion rate practicable. Ensure adequate hydration before administration. Monitor for signs and symptoms of thrombosis and assess blood viscosity in patients at risk for hyperviscosity.

Aseptic Meningitis Syndrome: Has been reported with use of IG and may occur more frequently in females. Conduct a thorough neurological exam on patients exhibiting signs and symptoms, to rule out other causes of meningitis. Discontinuing IG treatment has resulted in remission within several days without sequelae.

Hemolysis: GAMMAGARD LIQUID contains blood group antibodies, which may cause a positive direct antiglobulin reaction and hemolysis. Monitor patients for signs and symptoms of hemolysis and delayed hemolytic anemia and, if present, perform appropriate confirmatory lab testing.

Transfusion-Related Acute Lung Injury: Non-cardiogenic pulmonary edema has been reported with IV-administered IG, including GAMMAGARD LIQUID. Monitor patients for pulmonary adverse reactions. If suspected, perform appropriate tests for presence of anti-neutrophil and anti-HLA antibodies in both product and patient serum. May be managed using oxygen therapy with adequate ventilatory support.

Transmittable Infectious Agents: Because GAMMAGARD LIQUID is made from human plasma, it may carry a risk of transmitting infectious agents (e.g., viruses, other pathogens). No confirmed cases of viral transmission or variant Creutzfeldt-Jakob disease (vCJD) have been associated with GAMMAGARD LIQUID.

Interference with Lab Tests: False positive serological test results and certain assay readings, with the potential for misleading interpretation, may occur as the result of passively transferred antibodies.

Adverse Reactions

The most serious adverse reactions observed in clinical studies in PI was aseptic meningitis, and in MMN were pulmonary embolism and blurred vision.

The most common adverse reactions observed in ≥5% of patients were:

IV administration for PI: Headache, fatigue, pyrexia, nausea, chills, rigors, pain in extremity, diarrhea, migraine, dizziness, vomiting, cough, urticaria, asthma, pharyngolaryngeal pain, rash, arthralgia, myalgia, oedema peripheral, pruritus, and cardiac murmur.

Subcutaneous administration for PI: Infusion site (local) event (rash, erythema, edema, hemorrhage, and irritation), headache, fatigue, heart rate increased, pyrexia, abdominal pain upper, nausea, vomiting, asthma, blood pressure systolic increased, diarrhea, ear pain, aphthous stomatitis, migraine, oropharyngeal pain, and pain in extremity.

IV administration for MMN: Headache, chest discomfort, muscle spasms, muscular weakness, nausea, oropharyngeal pain, and pain in extremity.

IV administration for CIDP: Headache, pyrexia, anemia, leukopenia, neutropenia, illness, blood creatinine increased, dizziness, migraine, somnolence, tremor, nasal dryness, abdominal pain upper, vomiting, chills, nasopharyngitis, and pain in extremity.

Drug Interactions

Passive transfer of antibodies may transiently interfere with immune responses to live attenuated virus vaccines (e.g., measles, mumps, rubella, and varicella).

Please click for Full Prescribing Information.